

Spectroscopy and microchip laser operation of Tm, Ho:KYW crystals with different Ho concentrations

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Astro Ltd. The spectroscopic properties of Tm, Ho:KYW crystals with different Ho concentrations were investigated. The diode-pumped microchip laser operation of Tm (5 at.%), Ho (0.5 at.):KYW and Tm (5 at.%), Ho (1 at.):KYW was demonstrated. The highest, to our knowledge, output power of 480 mW with slope efficiency of 31% for CW Tm (5 at.%), Ho (0.5 at.):KYW microchip laser was obtained.

<http://dx.doi.org/10.1088/1612-202X/aa9972>

Keywords

absorption, double tungstates, holmium, microchip laser, thulium, up-conversion

References

- [1] Koch G J et al 2004 Coherent differential absorption lidar measurements of CO₂ Appl. Opt. 43 5092-9
- [2] Ghibaudo J B, Labandibar J Y, Marini A and Steinvall K O 1997 Water vapour and wind measurements by a two micron space lidar Proc. of the 9th Conf. on Coherent Laser Radar pp 7-10
- [3] Scholle K, Lamrini S, Kookmann P and Fuhrberg P 2010 2 μ laser sources and their possible applications Frontiers in Guided Wave Optics and Optoelectronics ed B Pal (Rijeka: InTech)
- [4] Bourdet G L and Lescroart G 1999 Theoretical modeling and design of a Tm,Ho:YLiF₄ microchip laser Appl. Opt. 38 3275-81
- [5] Izawa J, Nakajima H, Hara H and Arimoto Y 2000 A tunable and longitudinal mode oscillation of a Tm,Ho:YLF microchip laser using an external etalon Opt. Commun. 180 137-40
- [6] Bourdet G L and Muller R A 2000 Tm,Ho:YLF microchip laser under Ti:sapphire and diode pumping Appl. Phys. B 70 345-9
- [7] Yao B Q, Chen F, Meng P B, Zhang C H and Wang Y Z 2011 Diode pumped operation of Tm,Ho:YAP microchip laser Laser Phys. 21 674-6
- [8] Ju Y L, Wang Z G, Li Y F and Wang Y Z A 2008 Single-longitudinal-mode dual-wavelength cw Tm,Ho:GdVO₄ microchip laser Chin. Phys. Lett. 25 3250-2
- [9] Zhou R L, Ju Y L, Wu C T, Wang Z G and Wang Y Z 2010 A single-longitudinal-mode dual-wavelength CW Tm,Ho:GdVO₄ microchip laser Laser Phys. 20 1320-3
- [10] Yao B Q, Chen F, Wu C T, Wang Q, Li G, Zhang C H, Wang Y Z and Ju Y L 2011 Diode-end-pumped Tm,Ho:YVO₄ microchip laser at room temperature Laser Phys. 21 663-6
- [11] Storm M E and Rohrbach W W 1989 Single-longitudinal-mode lasing of Ho:Tm:YAG at 2.091 μ m Appl. Opt. 28 4965-7
- [12] Loiko P, Serres J M, Mateos X, Yumashev K, Kuleshov N, Petrov V, Griebner U, Aguiló M and Diaz F 2014 Microchip laser operation of Tm,Ho:KLu (WO₄)₂ crystal Opt. Express 22 27976-84
- [13] Kurilchik S, Gusakova N, Demesh M, Yasukevich A, Kisel V, Pavlyuk A and Kuleshov N 2016 Energy transfer in Tm,Ho:KYW crystal and diode-pumped microchip laser operation Opt. Express 24 6451-8

- [14] Lagatsky A A, Fusari F, Kurilchik S V, Kisel V E, Yasukevich A S, Kuleshov N V, Pavlyuk A A, Brown C T A and Sibbet W 2009 Optical spectroscopy and efficient continuous-wave operation near 2 μm for a Tm, Ho:KYW laser crystal Appl. Phys. B 97 321
- [15] Sani E, Toncelli A, Tonelli M, Coluccelli N, Galzerano G and Laporta P 2005 Appl. Phys. B 81 847-51
- [16] Han X, Fusari F, Serrano M D, Lagatsky A A, Cano-Torres J M, Brown C T A, Zaldo C and Sibbett W 2010 Continuous-wave laser operation of Tm and Ho co-doped NaY (WO₄)₂ and NaLu (WO₄)₂ crystals Opt. Express 18 5413-9
- [17] Walsh B M, Barnes N P and Di Bartolo B 2000 The temperature dependence of energy transfer between the Tm 3F₄ and Ho 5I₇ manifolds of Tm-sensitized Ho luminescence in YAG and YLF J. Lumin. 90 39-48
- [18] Bowman S R, Winings M J, Auyeung R C Y, Tucker J E, Searles S K and Feldman B J 1991 Laser and spectral properties of Cr,Tm,Ho:YAG at 2.1 μm IEEE J. Quantum Electron. 27 2142
- [19] Petrin R R, Jani M G, Powell R C and Kokta M 1992 Spectral dynamics of laser-pumped Y₃Al₅O₁₂:Tm, Ho lasers Opt. Mater. 1 111-24
- [20] Bigotta S, Toncelli A, Tonelli M, Cavalli E and Bovero E 2007 Spectroscopy and energy transfer parameters of Tm³⁺- and Ho³⁺-doped Ba₂NaNb₅O₁₅ Opt. Mater. 30 129-31
- [21] Serres J M, Loiko P, Mateos X, Jambunathan V, Yumashev K, Griebner U, Petrov V, Aguilo M and Diaz F 2015 Q-switching of a Tm,Ho:KLu (WO₄)₂ microchip laser by a graphene-based saturable absorber Laser Phys. Lett. 13 1-5
- [22] Pavlyuk A A, Vasiliev Ya V, Kharchenko L Yu and Kuznetsov F A 1992 Proc. of the APSAM-92 (Shanghai, China,) pp 164-71
- [23] Klevtsov P V, Kozeeva L P and Klevtsova R F 1968 Izv. Akad. Nauk. Neorg. Mat. 4 1147
- [24] Troshin A E, Kisel V E, Yasukevich A S, Kuleshov N V, Pavlyuk A A, Dunina E B and Kornienko A A 2007 Spectroscopy and laser properties of Tm³⁺:KY (WO₄)₂ crystal Appl. Phys. B 86 287-92
- [25] Carnal W T, Fields P R and Rajnak K 1968 Electronic energy levels in trivalent lanthanide aquo ions. I. Pr³⁺, Nd³⁺, Pm³⁺, Sm³⁺, Dy³⁺, Ho³⁺, Er³⁺, and Tm³⁺ J. Chem. Phys. 49 4424
- [26] Jambunathan V, Mateos X, Loiko P, Serres J M, Griebner U, Petrov V, Yumashev K, Aguilo M and Diaz F 2016 Growth, spectroscopy and laser operation of Ho:KY (WO₄)₂ J. Lumin. 179 50-8